

DRY SODIUM CHLORIDE AEROSOL IN REHABILITATION OF PATIENTS WITH COPD

Alina Chervinskaya¹, Irena Ponikowska², Robert Szafkowski², Jacek Chojnowski²

¹Clinical Research Rehabilitation Pulmonary Center, Hospital 122 of Federal Medical Biological Agency, St.Petersburg, Russian Federation

²Clinic of Balneology and Physical Medicine, Medical University in Bydgoszcz, Ciechocinek, Poland



Introduction

Management of COPD involves optimising medical therapy, commencing smoking cessation and participating in pulmonary rehabilitation (RT). Recently, there has been interest in ways of involving physical methods to RT.

Dry sodium chloride aerosol (DSCA) is the main acting factor of the speleotherapy (salt cave therapy) and halotherapy (therapy in a controlled air medium which saturated with dry salt aerosol). DSCA is characterized with physical properties, differing from those of the saline aerosols.

Our previous studies demonstrated that DSCA possessed anti-inflammatory activity in the respiratory tract, mucoregulating action. It enhances drainage of the bronchi, activates alveolar macrophages, improves biocenosis and local humoral immunity.

Aim of the study

The main objective was to estimate the efficacy of inhaled dry sodium chloride aerosol (DSCA) in rehabilitation therapy (RT) of patients with COPD.

Methods

It was double-blind placebo study. 72 patients (pts) with moderate and mild stage of COPD were recruited. They were randomized in 2 groups - interventional group (IG) (21 m, 18 f, 60.3±10.8 yrs) and control group (CG) (22 m, 11 f, 58.5±8.9 yrs). All patients received RT: daily procedures of chest massage, light radiation, physical exercises. Pts of IG were treated with the DSCA (45 min twice a day for 14 days). (Fig.1).




Figure #1

DSCA containing particles with size of 1-5 µm and level of mass concentration in the room of 10-15 mg/m³ (fig 2.) was produced by halogenerator GDA-01.17 (Halomed, Lithuania). CG received placebo (inhalations with room air) instead of DSCA.

Clinical, functional parameters and measures of health-related quality of life (HRQL) by SF-16 and LCQ (10 items) were estimated after RT procedures and in 3 months.

Modalities of dry sodium chloride aerosol at COPD rehabilitation programmes

Density range of dry sodium chloride aerosol 10-15 mg/m³

•Fractional composition of dry salt aerosol:
Respirable particles (1- 5 µm) – more, than 80%




Figure #2

Results

Improvements of clinical symptom scores were observed in the both groups after the course of RT (p<0.05), but in 3 months positive effect was noticed only in IG (before-13.8±5.4, after RT- 9.1±4.9, in 3 months -9.6±4.3, p<0.05). Measures of LCQ were changed significantly after RT only in pts of IG, received DSCA (35.2±5.2 and 52.4±6.3, p<0.05).

Positive changes of physical functioning measures (SF-16) were observed in IG and CG groups after RT, but they have been kept till 3 month only in IG.

Conclusions

Application of inhalations of DSCA on the background of the RT in pts with COPD renders to positive effect.

Bibliography

- Chervinskaya A. Halotherapy in controlled salt chamber microclimate for recovering medicine // Balneol. Pol. – 2007. – Vol. 49, N 2 (108). - P. 133-141.
- Chervinskaya A. V. Respiratory hygiene with the dry sodium chloride aerosol // 14th Annual Congress of the European Respiratory Society, Glasgow, September 2004: Abstract Book.- 2004. – Ref. 2514.
- Chervinskaya A.V. Effect of dry sodium chloride aerosol on the respiratory tract of tobacco smokers // The Europ. Respir. Journ.-Abstracts 16th ERS Annual Congress, Munich, Germany, September 2-6, 2006. – P. 106s-107s.
- Mokina N. A., Geppé N. A. Alternative methods at bronchial asthma of children // 14th Annual Congress of the European Respiratory Society, Glasgow, September 2004: Abstract Book.- 2004. – Ref. 1069.

22nd ERS Congress, Vienna, Austria
September 1-5 2012, P3549